

**METABOLIC RESPONSE OF ENVIRONMENTALLY ISOLATED
MICROORGANISMS TO INDUSTRIAL EFFLUENTS: USE OF A NEWLY
DESCRIBED CELL CULTURE ASSAY**

**Final Report
NASA/ASEE Summer Faculty Fellowship Program -1992
Johnson Space Center**

Prepared By: Robert N. Ferebee, Ph.D.
Academic Rank: Associate Professor
University and Department: University of Houston - Clear
Lake
Department of Biological
And Environmental Sciences
Houston, Texas 77058

NASA/JSC

Directorate: Space and Life Sciences
Division: Medical Sciences
Branch: Biomedical Operations and
Research

JSC Colleague: Duane L. Pierson, Ph.D.
Date Submitted: September 15, 1992
Contact Number:

**Metabolic Response of Environmentally Isolated Microorganisms To
Industrial Effluents; Use of a Newly Described Cell Culture Assay.**

**Robert N. Ferebee, Ph.D.
University of Houston-Clear Lake
June 16, 1992**

**Duane L. Pierson, ph.D.
Biomedical Research and Operations Division**

Abstract of Proposed Work Plan

An environmental application using a microtiter culture assay to measure the metabolic sensitivity of microorganisms to petrochemical effluents will be tested.

The Biomedical Operations and Research Branch at NASA JSC has recently developed a rapid and nondestructive method to measure cell growth and metabolism. Using a colorimetric procedure the uniquely modified assay allows the metabolic kinetics of prokaryotic and eukaryotic cells to be measured. Use of such an assay if adapted for the routine monitoring of waste products, process effluents, and environmentally hazardous substances may prove to be invaluable to the industrial community.

The microtiter method as described will be tested using microorganisms isolated from the Galveston Bay aquatic habitat. The microbial isolates will be identified prior to testing using the automated systems available at JSC. Sodium dodecyl sulfate (SDS), cadmium, and lead will provide control toxic chemicals. The toxicity of industrial effluent from two industrial sites will be tested.

An effort will be made to test the efficacy of this assay for measuring toxicity in a mixed culture community.

Metabolic Response of Environmentally Isolated Microorganisms to Industrial Effluents: Use of a Newly Described Cell Culture Assay.

Introduction:

Bioassays performed upon industrial process effluents continue to present problems related to time, difficulty in maintaining the animal species used, and the expense of such procedures. While this study does in no way suggest that bioassays using higher life forms be discontinued it does suggest there is a need to develop a rapid screening procedure that allows a daily monitoring of potentially toxic effluents.

Use of a recent modification of the tetrazolium reduction assay that allows simultaneous solubilization of the end product formazan indicated that the method could be used as a rapid screening assay for industrial wastes. This procedure not only allows one to detect changes in the physicochemical nature of effluents, but also provides a method to quickly screen for microorganisms that would be sensitive to toxic substances.

This research was designed to measure the effects that industrial effluents taken from three separate industrial process plants had on actively growing bacteria. The treated effluents from each of these industries are ultimately released into the Galveston Bay estuary system. Each of the industries were eager to cooperate in supplying samples for these studies.

Materials and Methods:

Test Organisms ; Nine separate bacterial genera were collected in grab samples taken from selected sites on Galveston Bay - Texas. The bacteria were identified to species using both " Vitek" (Vitek Systems Inc., Hazelwood, Mo.) and " Biolog" (Biolog Inc., Hayward, Ca.), computer assisted, automated systems. These environmental isolates were tested in the presence of control chemicals of known toxicity as well as both treated and untreated (raw) industrial effluents. Of the bacteria tested Staphylococcus hominis and Pseudomonas putida showed greatest promise as indicator species and were used in subsequent tests. These same organisms were obtained from stock culture collections (Chrisope Technologies Lake Charles, LA) ; the sensitivity of these bacteria were compared to that of the environmental isolates.

Chemicals; Chemicals for toxic controls were obtained from Mallinckrodt Chemical Works St. Louis, Mo.. Cadmium Chloride, Mercuric Sulfate, Lead Acetate, and Sodium Dodecyl Sulfate were all tested. Cadmium chloride gave the most consistent results with all the bacteria tested and was used for subsequent testing. Chemical for the tetrazolium assay were obtained from Sigma Chemical Co., St. Louis, Mo. Tetrazolium 1-(4,5- dimethylthiazol-2-yl)- 3,5 - diphenylformazan (MTT formazan), 2-phenazine methosulfate (PMS), and t-octylphenoxypolythoxyethanol (Triton X-100) were used in the assay for bacterial growth.

Culture media were obtained from BBL Microbiological Systems (Cockeysville, Md.). Bacteria were maintained on standard methods agar while the media used for microtiter culture was R2A broth.

Tetrazolium Assays: Testing for the bacterial activity was measured by reduction of MTT spectrophotometrically at 390nm minus 650nm. Test bacteria were removed from 18 hour cultures by sterile cotton swabs and suspended in filter sterilized deionized water. Cell suspensions were adjusted to OD 590 nm = 0.6 cell numbers. This density gave 2×10^9 cells/ml. From this suspension 25 ul were added to 50 ul of double strength R2A broth in 96 well flat bottomed microtiter culture plates. All wells except cell controls received 50 ul known toxicants or 50 ul treated or raw effluent. Indicator chemicals were finally added as follows: 15 ul of 1.2 mM MTT, 10 ul of 0.06 mM PMS, and 25 ul of 160 mM Triton X-100. All controls and tests were set as replicates of four wells. Cultures were maintained at 30C and continually monitored at 590nm - 650nm for a maximum of two hours. The microtiter cultures were read using a microtiter plate reader (Molecular Device Co. Menlo Park, CA).

Results:

Each of the environmental bacteria as well as those obtained from stock collections exhibited an expected response when toxic chemicals were added to the cultures. There was a progressive inhibition of growth as the concentrations of the chemicals were increased, (Fig 1 & 2). The cadmium chloride gave the most consistent response and was used as the toxicant control (Fig 1&2). None of the chemicals tested caused any reduction of MTT; these represented chemical controls. When the bacteria selected for indicators were tested with industrial effluents only a single sample gave a significant change in bacterial growth and MTT reduction (figure 1 & 2). The sample identified as effluent (2R) did indicate toxicity; further testing of this effluent showed a pH of 2.8 could account for the reduction in growth. When this sample was buffered to pH 7.0 ± 0.5 the toxicity was reversed. Staphylococcus hominis cultures were not effected by any of the effluent samples; while the Pseudomonas putida indicated changes that were minimum.

Discussion:

This new procedure for rapid reduction of tetrazolium allows for a rapid assay with good sensitivity and processing of large numbers of samples. The procedure offers a simple procedure that can easily be adapted for industrial operations. As testing continues there will probably be other microbial indicators identified that may be as good for screening as the ones used in this study. It is deemed somewhat important to choose environmental isolates that are part of the resident population. Many of these resident organisms represent members of the natural

bioremediation/biodegradation community that ecosystems have become dependent upon for recycling, etc;. As is suggested by this study it may be beneficial to name more than a single indicator genera or even different species. Some of the preliminary results suggested that some of the chemicals will enhance the growth of certain bacteria. This was true in the case of sodium dodecyl sulfate (SDS) as a control toxicant. This type of activity indicates the eutrophic activity of many chemicals and only supports the value of this procedure in the testing of effluents.

TABLE 1: BACTERIAL ISOLATES COLLECTED FROM GALVESTON BAY - TEXAS

Acinetobacter calcaeticus

Aeromonas veronii

Alcaligenes faecalis

Bacillus coagulans

Enterobacter cloacae

Pseudomonas putida

Serratia marcesens

Staphylococcus hominis

Vibrio metschnikovii





Figure 1: Staphylococcus hominis
Growth in Industrial Effluent



**Figure 2: *Pseudomonas putida*
Growth in Industrial Effluent**

LITERATURE REVIEWED

1. Bartlett, R. C., M. Mazens, and B. Greenfield. 1976. Acceleration of tetrazolium reduction by bacteria. *J. Clin. Microbiol.* 3(3): 327-329.
2. Bochner, B.A., and M.A. Savageau. 1977. Generalized Indicator Plate for Genetic, Metabolic, and Taxonomic Studies with Microorganisms. *Applied and Environmental Microbiology*. Vol.33 No.2, p. 434-444.
3. Denizot, F. and R. Lange. 1986. Rapid colorimetric assay for cell growth and survival: Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J. Immunological Methods* (89): 271-277.
4. Findlay, R. H. , M. B. Trexler, J. B. Guckert, and D. C. White. 1990. Laboratory study of disturbance in marine sediments: response of a microbial community. *Marine Ecology Prog. Ser.* Vol. 62: 121-133.
5. Garland, J. L., and A. L. Mills. 1991. Classification and Characterization of Heterotrophic Microbial Communities on the Basis of Patterns of Community- Level Sole-Carbon-Source Utilization. *Applied and Environmental Microbiology*. Vol. 57, No.8. 2351-2359.
6. Gillan, F. T. and R.W. Hogg. 1984. A Method for the estimation of Bacterial biomass and Community Structure in Mangrove-Associated sediments. *Journal of Microbiological Methods*. 2:275-293.
7. Jeffrey, W. H., and J.H. Paul. 1986. Activity of an attached and free-living *Vibrio* sp. as measured by thymidine incorporation, p-iodonitrotetrazolium reduction, and ATP/DNA ratios. *Appl. Environ. Microbiol.* 51(1):150-156.
8. Kasugai, S. N. Hasegawa, and H. Ogura. 1990. A simple in vitro cytotoxicity test using the MTT (3-(4,5)-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) colorimetric assay: Analysis of eugenol toxicity on dental pulp cells (RPC-C2A). *Japan J. Pharmacol.* 52:59-100.
9. Karmer, V. C., D. M. Calabrese, and K. W. Nickerson. 1980. Growth of *Enterobacter cloacae* in the presence of 25 μ g sodium dodecyl sulfate. *Appl. Environ. Microbiol.* 40: 711-713.
10. Mosmann, T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods*. 65: 55-63.

11. Page' M., N. Bejaoui, B. Cinq-Mars and P. Lemieux. 1988. Optimization of the Tetrazolium - Based Colorimetric Assay for the Measurement of Cell Number and Cytotoxicity. Vol. 10. No. 7 p.785-793.
12. Roszak, D. B. and R. R. Colwell. 1987. Survival Strategies of Bacteria in the Natural Environment. Microbiological Reviews.Vol. 51. No. 3. p.365-379.
13. Schnaitman, C. A. 1971. Effect of Ethylenediaminetetracetic acid, triton X-100, and lysozyme on the morphology and chemical composition of isolated cell walls of Escherichia coli. J. Bacteriology 108: 553-563.
14. Stowe, R. A.,D. W. Koenig,S.K. Mishra, and D. L. Pierson.1992. In preparation. A Nondestructive Tetrazolium Microtiter Assay for Measurement of Cell Metabolism and Growth. Biomedical Operations and Research Branch National Aeronautics and Space Administration, Lyndon B. Johnson Space Center, Mail Code SD4 Houston, Tx. 77058.
15. Weber, C. A. et.al. 1988. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Eceiving Waters to Marine and Estuarine Organisms. Biological Methods Branch Environmental Monitoring and Support Laboratory - Cincinnati , U. S. Environmental Protection Agency Cincinnati, Ohio 45268.